Application No. 10/765,678 SD-7463

# BEST AVAILABLE COPY

### AMENDMENTS TO THE CLAIMS

- Please amend the claims as follows:
- 1. (Currently Amended) An adueous decontamination formulation for use in disinfection and sterilization, said formulation comprising consisting essentially of:
  - a reactive compound selected from the group consisting of nucleophilic compounds and oxidizing compounds;
  - a water-soluble bleaching activator selected from the group consisting of shortchained-organic compounds that contain an ester bend, ethylene glycol
    diacetate, propy ene glycol monomethyl ether acetate, methyl acetate,
    diethylene glycol monoethyl ether acetate, glycerol acetate (monoacetin),
    glycerol diacetate (diacetin), glycerol triacetate (triacetin), acetylcholine
    chloride, 4-cyanobenzoic acid, propylene glycol diacetate, and
    combinations thereof;

an inorganic base; and

water.

wherein said formulation does not comprise a solubilizing compound selected from the group consisting of a cationic surfactant, a cationic hydrotrope, and a fatty alcohol comprising 8 20 carbon atoms.

- 2. (Currently Amended) The formulation of claim 1, comprising consisting essentially of (by weight percentage):
  - 0.5-60 % reactive corppound;
  - 1-10 % bleaching activator;
  - 3-30% inorganic base
  - 0 10% ethanol;
  - 0 20 % freeze point depressant; and water (remainder).

Application No. 10/765,678 SD-7463

### **BEST AVAILABLE COPY**

- 3. (Currently Amended) The formulation of claim 2, wherein:
  said reactive compound comprises hydrogen peroxide;
  said bleaching activator comprises glycerol diacetate or propylene glycol
  diacetate; and
  said inorganic base comprises potassium acetate; and
  said freeze point depressant comprises propylene glycol.
- 4. (currently amended) The formulation of claim 1, further comprising An aqueous decontamination formulation for use in disinfection and sterilization, said formulation consisting essentially of:
  - a reactive compound selected from the group consisting of nucleophilic compounds and oxidizing compounds:
  - a water-soluble bleaching activator selected from the group consisting of
    ethylene glycol diacetate, propylene glycol monomethyl ether acetate,
    methyl acetate, diethylene glycol monoethyl ether acetate, glycerol
    acetate (monoacetin), glycerol diacetate (diacetin), glycerol triacetate
    (triacetin), acetylcholine chloride, 4-cyanobenzoic acid, propylene glycol
    diacetate, and combinations thereof:

#### an inorganic base;

#### water; and

- one or more sorber t additives selected from the group consisting of sodium carbonate, sodium bicarbonate, potassium carbonate, potassium bicarbonate, calcium carbonate, potassium silicate, precipitated silicates, percarbonates, amorphous silica, fumed silica, sodium citrate, dendritic salt (sea salt), citric acid, polyethylene glycol, PEG 8000, urea, and polyols.
- 5. (original) The formulation of claim 4, wherein said sorbent additive comprises one or more polyol compounds selected from the group consisting of sorbitol, mannitol, hydrogenated starch hydrolysates (HSH), maltitol, zylitol, lactitol monohydrate, anhydrous isomalt, erythritol, and polydextrose.

+15058442829

Application No. 10/765,678 SD-7463

## BEST AVAILABLE COPY

- 6. (Currently Amended) An aqueous decontamination formulation for use in neutralization of a toxant, said formulation comprising consisting essentially of (by weight percentage):
  - 0.5-60 % hydrogen perloxide;
  - 1-10 % glycerol diacetate or propylene glycol diacetate;
  - 3-10% potassium carbbnate;
  - 0-10% ethanol; and

water (remainder);

wherein said formulation does not comprise a solubilizing compound selected from the group consisting of a cationic surfactant, a cationic hydrotrope, and a fatty alcohol comprising 8-20 carbon atoms.

- 7. (Currently Amended) An aqueous decontamination formulation for use in neutralization of a toxant, said formulation comprising consisting essentially of (by weight percentage):
  - 0.5-60 % hydrogen peroxide;
  - 1-10 % glycerol diacetate or propylene glycol diacetate;
  - 0-5%-benzalkonium-dhloride;
  - 5-30% potassium acetate;
  - 0.20% proplyene glydol; and
  - water (remainder);
  - wherein said fermulation-comprises no amount of a carbonate or bicarbonate salt.
- 8-28. CANCELLED
- 29. (Previously Presented) The formulation of claim 1, wherein the inorganic base is selected from the group consisting of potassium carbonate, potassium bicarbonate,

Application No. 10/765,678 SD-7463

## BEST AVAILABLE COPY

potassium hydroxide, potassium sulfate, potassium phosphate (dibasic or tribasic), potassium borate, potassium tetraborate, potassium acetate, sodium carbonate, sodium bicarbonate, sodium hydroxide, sodium sulfate, sodium phosphate (dibasic or tribasic), sodium borate, sodium acetate, ammonium carbonate, ammonium bicarbonate, ammonium hydroxide, ammonium sulfate, ammonium phosphate (dibasic or tribasic), ammonium borate, ammonium acetate, calcium carbonate, calcium bicarbonate, calcium hydroxide, calcium sulfate, calcium phosphate (dibasic or tribasic), calcium borate, calcium acetate, magnesium carbonate, magnesium bicarbonate, magnesium hydroxide, magnesium sulfate, magnesium phosphate (dibasic or tribasic), magnesium borate, magnesium acetate, sodium percarbonate, ammonium hydrogen bicarbonate and lithium bicarbonate, and combinations thereof.